

Notes on matrix inequalities and positive multilinear mappings

Leila Nasiri

Department Mathematics, Lorestan University, Iran

(Communicated by Abbas Najati)

Abstract

In this paper, our aim is to prove some matrix inequalities involving arbitrary matrix means and positive multilinear mappings. For example, it is shown that for Hermitian matrices A_i, B_i such that $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$),

$$\Phi^2(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \leq (K(u^k))^2 \Phi^2(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k),$$

where σ_1 and σ_2 are two arbitrary matrix means between the arithmetic and harmonic means, $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_l(\mathbb{C})$ is a positive unital multilinear mapping, $u = \frac{M}{m}$ and $K(u) = \frac{(1+u)^2}{4u}$. We also give the obtained results for the adjoint and the dual of an arbitrary matrix mean.

Keywords: Matrix mean, Positive multilinear mapping, Positive matrix
2020 MSC: 47A63, 47A30, 47A64

1 Introduction

Let $\mathcal{M}_n(\mathbb{C})$ denote the C^* -algebra of all $n \times n$ matrices with entries in the complex field \mathbb{C} with the identity I . For a Hermitian matrix $A \in \mathcal{M}_n(\mathbb{C})$, we write $A \geq 0$ if all its eigenvalues are non-negative. We also write $A > 0$, if it be invertible, moreover the before condition. For two Hermitian matrices $A, B \in \mathcal{M}_n(\mathbb{C})$, we use the notation $A \leq B$ (or $A > B$) to mean that $B - A \geq 0$ (or $B - A > 0$). A mapping $\Phi : \mathcal{M}_n^k(\mathbb{C}) = \mathcal{M}_n(\mathbb{C}) \times \dots \times \mathcal{M}_n(\mathbb{C}) \rightarrow \mathcal{M}_l(\mathbb{C})$ is said to be multilinear whenever it is linear in each of its variable and also is called positive if $A_i \geq 0$ for $i = 1, \dots, k$ implies that $\Phi(A_1, \dots, A_k) \geq 0$. Moreover, Φ is called unital if $\Phi(I, \dots, I) = I$. If $A \in \mathcal{M}_n(\mathbb{C})$ is a Hermitian matrix, the Gelfand map $f(t) \mapsto f(A)$ is an isometrically $*$ -isomorphism between the C^* -algebra $C(\text{sp}(A))$ of continuous functions on the spectrum $\text{sp}(A)$ of the Hermitian matrix A and the C^* -algebra generated by I and A . If $f, g \in C(\text{sp}(A))$, then $f(t) \geq g(t)$ ($t \in \text{sp}(A)$) implies that $f(A) \geq g(A)$. For $J \subset \mathbb{R}$, a real valued continuous function $f : J \rightarrow \mathbb{R}$ is called matrix monotone if $A \leq B$ implies that $f(A) \leq f(B)$ for all Hermitian matrices A and B whose eigenvalues are in J . Kubo and Ando [1] defined the matrix mean σ for pairs of positive definite matrices A and B and the nonnegative matrix monotone function $f : (0, \infty) \rightarrow (0, \infty)$ with $f(1) = 1$

$$A\sigma B = A^{\frac{1}{2}}f(A^{-\frac{1}{2}}BA^{-\frac{1}{2}})A^{\frac{1}{2}}$$

where f is the representing function for σ .

For two positive definite matrices $A, B \in \mathcal{M}_n(\mathbb{C})$, the Löwner–Heinz inequality states that, if $A \leq B$, then

$$A^p \leq B^p, \quad (0 \leq p \leq 1). \tag{1.1}$$

Email address: leilanasiri468@gmail.com (Leila Nasiri)

In general (1.1) is not true for $p > 1$. The authors [8] proved that if A and B are two positive definite matrices such that $0 < m \leq A, B \leq M$, then

$$\Phi^2(A\sigma_1B) \leq \left(\frac{(M+m)^2}{4Mm}\right)^2 \Phi^2(A\sigma_2B), \tag{1.2}$$

where Φ is a positive unital linear mapping and σ_1, σ_2 are two arbitrary matrix means between the harmonic and arithmetic matrix means. The authors in [6] generalized the inequality (1.2) for $p > 0$ as follows:

$$\Phi^p(A\sigma_1B) \leq \left(\frac{(M+m)^2}{4Mm}\right)^p \Phi^p(A\sigma_2B), \tag{1.3}$$

where $0 < m \leq A, B \leq M$, Φ is a positive unital linear mapping, σ_1, σ_2 are two arbitrary matrix means between the harmonic and arithmetic matrix means. For more information on the above inequalities see [5, 10, 12, 13, 14, 15, 7] and references therein. In this paper, we intend to extend the inequality (1.3) for multilinear positive mappings. Then, we shall generalize the derived results for the adjoint and the dual of an arbitrary matrix mean.

2 Main results

2.1 Some inequalities for arbitrary matrix means

This section is started by recalling several well known lemmas.

Lemma 2.1. [4] Suppose that $A_i \in \mathcal{M}_n(\mathbb{C})$ ($i = 1, \dots, k$) be positive definite. Then for every unital multilinear mapping $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_l(\mathbb{C})$

$$\Phi(A_1, \dots, A_k)^{-1} \leq \Phi(A_1^{-1}, \dots, A_k^{-1}).$$

Lemma 2.2. [1, 2, 3] Let $A, B \in \mathcal{M}_n(\mathbb{C})$ be positive definite and $\alpha > 0$. Then

- (i) $\|AB\| \leq \frac{1}{4}\|A+B\|^2$.
- (ii) $\|A^\alpha + B^\alpha\| \leq \|(A+B)^\alpha\|$ for $\alpha \geq 1$.
- (iii) $A \leq \alpha B$ if and only if $\|A^{\frac{1}{2}}B^{-\frac{1}{2}}\| \leq \alpha^{\frac{1}{2}}$.

Lemma 2.3. [9] Let $X \in \mathcal{M}_n(\mathbb{C})$. Then $\|X\| \leq t$ if and only if

$$\begin{pmatrix} tI & X \\ X^* & tI \end{pmatrix} \geq 0.$$

Lemma 2.4. [13] Let f be a strictly positive convex twice differential function on the interval $[m, M]$ with $0 < m < M$ and let $C_i \in \mathcal{M}_n(\mathbb{C})$ such that $\sum_{i=1}^k C_i^* C_i = I$. If $A_i \in \mathcal{M}_n(\mathbb{C})$ such that $0 < m \leq A_i \leq M$ ($i = 1, \dots, k$), then

$$\sum_{i=1}^k C_i^* f(A_i) C_i \leq a_f \sum_{i=1}^k C_i^* A_i C_i + b_f I \leq \alpha f \left(\sum_{i=1}^k C_i^* A_i C_i \right),$$

where $a_f = \frac{f(M)-f(m)}{M-m}$, $b_f = \frac{Mf(m)-mf(M)}{M-m}$ and $\alpha = \max_{m \leq t \leq M} \left\{ \frac{a_f t + b_f}{f(t)} \right\}$.

First, we prove following efficient lemma.

Lemma 2.5. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ be such that $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$), for some scalars $0 < m < M$. Then, for every positive unital multilinear mapping $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_l(\mathbb{C})$,

$$\Phi(A_1, \dots, A_k) + M^k m^k \Phi(A_1^{-1}, \dots, A_k^{-1}) \leq M^k + m^k.$$

Proof . It follows the spectral decomposition of $A_i \in \mathcal{M}_n(\mathbb{C})$ that $A_i = \sum_{j=1}^n \lambda_{ij} Q_{ij}$ ($i = 1, \dots, k$) such that $\sum_{j=1}^n Q_{ij} = I$. Putting

$$C(j_1, \dots, j_k) = (\Phi(Q_{1j_1}, \dots, Q_{kj_k}))^{\frac{1}{2}}$$

such that

$$\sum_{j_1=1}^n \sum_{j_2=1}^n \cdots \sum_{j_k=1}^n C^*(j_1, \dots, j_k) C(j_1, \dots, j_k) = I.$$

One can get

$$\begin{aligned} & \Phi(A_1^{-1}, \dots, A_k^{-1}) \\ & \leq \Phi \left(\sum_{j=1}^n \lambda_{1j}^{-1} Q_{1j}, \dots, \sum_{j=1}^n \lambda_{kj}^{-1} Q_{kj} \right) \quad (\text{by the convexity of } f(t) = t^{-1}) \\ & = \sum_{j_1=1}^n \sum_{j_2=1}^n \cdots \sum_{j_k=1}^n \lambda_{1j_1}^{-1} \lambda_{2j_2}^{-1} \cdots \lambda_{kj_k}^{-1} \Phi(Q_{1j_1}, \dots, Q_{kj_k}) \quad (\text{by the multilinearity of } \Phi) \\ & \leq a_1 \sum_{j_1=1}^n \sum_{j_2=1}^n \cdots \sum_{j_k=1}^n C(j_1, \dots, j_k) \lambda_{1j_1} \lambda_{2j_2} \cdots \lambda_{kj_k} C(j_1, \dots, j_k) + b_1 I \\ & \hspace{15em} (\text{by Lemma 2.4}) \\ & = a_1 \sum_{j_1=1}^n \sum_{j_2=1}^n \cdots \sum_{j_k=1}^n \lambda_{1j_1} \lambda_{2j_2} \cdots \lambda_{kj_k} \Phi(Q_{1j_1}, \dots, Q_{kj_k}) + b_1 I \\ & = a_1 \Phi(A_1, A_2, \dots, A_k) + b_1 I, \end{aligned}$$

where $a_1 = \frac{-1}{M^k m^k}$ and $b_1 = \frac{M^k + m^k}{M^k m^k}$. Therefore,

$$\Phi(A_1, \dots, A_k) + M^k m^k \Phi(A_1^{-1}, \dots, A_k^{-1}) \leq M^k + m^k.$$

This proves the desired inequality. \square

By applying Lemma 2.5, we have the following result:

Lemma 2.6. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ be such that $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$), for some scalars $0 < m < M$ and $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_l(\mathbb{C})$ be a positive unital multilinear mapping, then

$$\Phi(A_1 \nabla B_1, \dots, A_k \nabla B_k) + M^k m^k \Phi(A_1^{-1} \nabla B_1^{-1}, \dots, A_k^{-1} \nabla B_k^{-1}) \leq (M^k + m^k). \tag{2.1}$$

Proof . Since Φ is multilinear and by Lemma 2.5, it follows that

$$\begin{aligned} & \Phi(A_1 \nabla B_1, \dots, A_k \nabla B_k) + M^k m^k \Phi(A_1^{-1} \nabla B_1^{-1}, \dots, A_k^{-1} \nabla B_k^{-1}) \\ & = \left(\frac{1}{2}\right)^k \left[\Phi(A_1, \dots, A_k) + M^k m^k \Phi(A_1^{-1}, \dots, A_k^{-1}) + \Phi(B_1, \dots, B_k) + M^k m^k \Phi(B_1^{-1}, \dots, B_k^{-1}) \right] \\ & \leq (M^k + m^k). \end{aligned}$$

\square

Remark 2.7. If we choose $k = 1$, in the inequality (2.1), then we get

$$\Phi(A \nabla B) + M m \Phi(A^{-1} \nabla B^{-1}) \leq M + m.$$

Lemma 2.8. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ be such that $0 < A_i \leq B_i$ ($i = 1, \dots, k$) and $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_l(\mathbb{C})$ be any positive unital multilinear mapping. Then

$$\Phi(A_1, \dots, A_k) \leq \Phi(B_1, \dots, B_k). \tag{2.2}$$

Proof . Since $0 < A_i \leq B_i$ ($i = 1, \dots, k$), and in result $B_i - A_i \geq 0$. Thus, we can write

$$\begin{aligned} \Phi(A_1, A_2, \dots, A_k) & \leq \Phi(B_1, A_2, \dots, A_k) \\ & \leq \Phi(B_1, B_2, \dots, A_k) \\ & \leq \Phi(B_1, B_2, \dots, B_k). \end{aligned}$$

\square

Theorem 2.9. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ be such that $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$), for some scalars $0 < m < M$ and let σ_1 and σ_2 be two arbitrary matrix means between the arithmetic and harmonic means. Then, for every positive unital multilinear mapping $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_l(\mathbb{C})$,

$$\Phi^2(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \leq (K(u^k))^2 \Phi^2(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k), \tag{2.3}$$

where $u = \frac{M}{m}$ and $K(u) = \frac{(1+u)^2}{4u}$.

Proof . By our assumption and using Lemma 2.6, we have

$$\begin{aligned} & \Phi(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) + M^k m^k \Phi((A_1\sigma_2B_1)^{-1}, \dots, (A_k\sigma_2B_k)^{-1}) \\ & \leq \Phi(A_1\nabla B_1, \dots, A_k\nabla B_k) + M^k m^k \Phi((A_1!B_1)^{-1}, \dots, (A_k!B_k)^{-1}) \\ & = \Phi(A_1\nabla B_1, \dots, A_k\nabla B_k) + M^k m^k \Phi(A_1^{-1}\nabla B_1^{-1}, \dots, A_k^{-1}\nabla B_k^{-1}) \\ & \leq (M^k + m^k). \end{aligned} \tag{2.4}$$

Applying Lemma 2.2(i), Lemma 2.1, respectively, and utilizing (2.4), we have

$$\begin{aligned} & \left\| \Phi(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) M^k m^k \Phi^{-1}(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) \right\| \\ & \leq \frac{1}{4} \left\| \Phi(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) + M^k m^k \Phi^{-1}(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) \right\|^2 \\ & \leq \frac{1}{4} \left\| \Phi(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) + M^k m^k \Phi((A_1\sigma_2B_1)^{-1}, \dots, (A_k\sigma_2B_k)^{-1}) \right\|^2 \\ & \leq \frac{(M^k + m^k)^2}{4}, \end{aligned}$$

which completes the proof. \square

Remark 2.10. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ be such that $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$), let σ_1 and σ_2 be two arbitrary matrix means between the arithmetic and harmonic means and $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_l(\mathbb{C})$ be a positive unital multilinear mapping. Then,

$$\Phi^p(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \leq (K(u^k))^p \Phi^p(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k), \quad 0 \leq p \leq 2 \tag{2.5}$$

where $K(u) = \frac{(1+u)^2}{4u}$ and $u = \frac{M}{m}$.

Note that for $0 \leq p \leq 2$, it follows $0 \leq \frac{p}{2} \leq 1$. So, by the (2.3) and the Lowner-Heinz inequality, we have (2.5).

The following theorem is an extension of inequality (2.5) for $p \geq 2$.

Theorem 2.11. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ such that $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$) and let σ_1 and σ_2 be two arbitrary matrix means between the arithmetic and harmonic means. Then for every positive unital multilinear mapping $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_l(\mathbb{C})$ and $p \geq 2$,

$$\Phi^p(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \leq (K(u^k))^p \Phi^p(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k), \tag{2.6}$$

where $u = \frac{M}{m}$ and $K(u) = \frac{(1+u)^2}{4u}$.

Proof . By Lemma 2.2(i) and (ii), Lemma 2.1 and (2.4), respectively, one can get

$$\begin{aligned} & \left\| \Phi^{\frac{p}{2}}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) M^{\frac{pk}{2}} m^{\frac{pk}{2}} \Phi^{-\frac{p}{2}}(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) \right\| \\ & \leq \frac{1}{4} \left\| \Phi^{\frac{p}{2}}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) + M^{\frac{pk}{2}} m^{\frac{pk}{2}} \Phi^{-\frac{p}{2}}(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) \right\|^2 \\ & \leq \frac{1}{4} \left\| (\Phi(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) + M^k m^k \Phi((A_1\sigma_2B_1)^{-1}, \dots, (A_k\sigma_2B_k)^{-1}))^{\frac{p}{2}} \right\|^2 \\ & = \frac{1}{4} \left\| \Phi(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) + M^k m^k \Phi((A_1\sigma_2B_1)^{-1}, \dots, (A_k\sigma_2B_k)^{-1}) \right\|^p \\ & \leq \frac{(M^k + m^k)^p}{4}. \end{aligned}$$

This shows (2.6). \square

Remark 2.12. If we take $k = 1$, the inequality (2.6) reduces the inequality (1.3). So, (2.6) is an extension of (1.3).

Now, we prove the following useful result to present the another extension of inequality (2.5) for $p \geq 4$.

Lemma 2.13. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ be such that $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$), for some scalars $0 < m < M$ and let σ_1 and σ_2 be two arbitrary matrix means between the arithmetic and harmonic means. Then, for every positive unital multilinear mapping $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_l(\mathbb{C})$, we have

$$\Phi^2(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) + M^{2k}m^{2k}\Phi^{-2}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \leq (M^{2k} + m^{2k}). \tag{2.7}$$

Proof . From $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$) and (2.2), it is clear that

$$\Phi(mI, \dots, mI) = m^k \leq \Phi(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \leq \Phi(MI, \dots, MI) = M^k. \tag{2.8}$$

We know that for every matrix T such that $0 < m \leq T \leq M$,

$$M^2m^2T^{-2} + T^2 \leq M^2 + m^2.$$

If we apply the same property for (2.8), (2.7) deduces. \square

In below, we obtain the another extension of (2.5) for $p \geq 4$.

Theorem 2.14. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ such that $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$) and let σ_1 and σ_2 be two arbitrary matrix means between the arithmetic and harmonic means. Then for every positive unital multilinear mapping $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_l(\mathbb{C})$ and $p \geq 4$,

$$\Phi^p(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \leq \left(\frac{K(u^k)(M^{2k} + m^{2k})}{4^{\frac{p}{2}}M^km^k} \right)^p \Phi^p(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k), \tag{2.9}$$

where $u = \frac{M}{m}$ and $K(u) = \frac{(1+u)^2}{4u}$.

Proof . By(2.3) and (2.7), we get

$$\begin{aligned} & K(u^k)\Phi^2(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) + \frac{M^{2k}m^{2k}}{K(u^k)}\Phi^{-2}(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) \\ & \leq K(u^k)\Phi^2(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) + M^{2k}m^{2k}K(u^k)\Phi^{-2}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \\ & \leq K(u^k)(M^{2k} + m^{2k}). \end{aligned} \tag{2.10}$$

Applying Lemma 2.2(i) and (ii), together with (2.10), we derive

$$\begin{aligned} & \left\| \Phi^{\frac{p}{2}}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k)M^{\frac{pk}{2}}m^{\frac{pk}{2}}\Phi^{-\frac{p}{2}}(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) \right\| \\ & \leq \frac{1}{4} \left\| \left[K^{\frac{p}{4}}(u^k)\Phi^{\frac{p}{2}}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) + \left(\frac{M^{2k}m^{2k}}{K(u^k)} \right)^{\frac{p}{4}}\Phi^{-\frac{p}{2}}(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) \right] \right\|^2 \\ & \leq \frac{1}{4} \left\| \left[K(u^k)\Phi^2(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) + \frac{M^{2k}m^{2k}}{K(u^k)}\Phi^{-2}(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) \right]^{\frac{p}{4}} \right\|^2 \\ & = \frac{1}{4} \left\| \left[K(u^k)\Phi^2(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) + \frac{M^{2k}m^{2k}}{K(u^k)}\Phi^{-2}(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) \right] \right\|^{\frac{p}{2}} \\ & \leq \frac{(K(u^k)(M^{2k} + m^{2k}))^{\frac{p}{2}}}{4}, \end{aligned}$$

which leads to the desired result. \square

Theorem 2.15. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ be such that $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$) with $0 < m < M$ and let σ_1 and σ_2 be two arbitrary matrix means between the arithmetic and harmonic means. Then, for every positive unital multilinear mapping $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_l(\mathbb{C})$ and $1 \leq \alpha \leq 2$,

$$\Phi^p(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \leq \frac{(K^{\frac{\alpha}{2}}(u^k)(M^{k\alpha} + m^{k\alpha}))^{\frac{2p}{\alpha}}}{16M^{kp}m^{kp}}\Phi^p(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k), \quad p \geq 2\alpha, \tag{2.11}$$

where $u = \frac{M}{m}$ and $K(u) = \frac{(1+u)^2}{4u}$ is the Kantorovich constant.

Proof . For $1 \leq \alpha \leq 2$, the (2.3) and the Löwner–Heinz inequality ensures us that

$$\Phi^\alpha(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) \leq (K(u^k))^\alpha \Phi^\alpha(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k),$$

which is equivalent to

$$\Phi^{-\alpha}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \leq (K(u^k))^\alpha \Phi^{-\alpha}(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k). \tag{2.12}$$

Making use of our assumptions, one can easily prove that

$$\Phi^\alpha(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) + M^{k\alpha}m^{k\alpha}\Phi^{-\alpha}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \leq M^{k\alpha} + m^{k\alpha}. \tag{2.13}$$

To prove (2.11), from Lemma 2.2(i) and (ii) and inequalities (2.12) and (2.13), we have

$$\begin{aligned} & \left\| M^{\frac{kp}{2}} m^{\frac{kp}{2}} \Phi^{\frac{p}{2}}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \Phi^{-\frac{p}{2}}(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) \right\| \\ & \leq \frac{1}{4} \left\| K^{-\frac{p}{4}}(u^k) M^{\frac{kp}{2}} m^{\frac{kp}{2}} \Phi^{-\frac{p}{2}}(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) + K^{\frac{p}{4}}(u^k) \Phi^{\frac{p}{2}}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \right\|^2 \\ & \leq \frac{1}{4} \left\| \left(K^{-\frac{\alpha}{2}}(u^k) M^{k\alpha} m^{k\alpha} \Phi^{-\alpha}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) + K^{\frac{\alpha}{2}}(u^k) \Phi^\alpha(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) \right)^{\frac{p}{2\alpha}} \right\|^2 \\ & = \frac{1}{4} \left\| K^{-\frac{\alpha}{2}}(u^k) M^{k\alpha} m^{k\alpha} \Phi^{-\alpha}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) + K^{\frac{\alpha}{2}}(u^k) \Phi^\alpha(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) \right\|^{\frac{p}{\alpha}} \\ & \leq \frac{K^{\frac{p}{2}}(u^k)(M^{k\alpha} + m^{k\alpha})^{\frac{p}{\alpha}}}{4}, \end{aligned}$$

This is equivalent to

$$\Phi^p(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) \leq \frac{(K^{\frac{\alpha}{2}}(u^k)(M^{k\alpha} + m^{k\alpha}))^{\frac{2p}{\alpha}}}{16M^{kp}m^{kp}} \Phi^p(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k)$$

This shows the desired inequality. \square

Remark 2.16. Choosing $\alpha = 1$ and $\alpha = 2$, in the inequality (2.11), we get to the inequalities (2.6) and (2.9), respectively.

Theorem 2.17. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ be such that $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$), when $0 < m < M$ and let σ_1 and σ_2 be two arbitrary matrix means between the arithmetic and harmonic means. Then, for every positive unital multilinear mapping $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_1(\mathbb{C})$,

$$\|\gamma_2 + \gamma_1\| \leq 2K^p(u^k), \quad p > 0,$$

where

$$\gamma_1 = \Phi^{\frac{p}{2}}(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k) \Phi^{-\frac{p}{2}}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k),$$

$$\gamma_2 = \Phi^{-\frac{p}{2}}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \Phi^{\frac{p}{2}}(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k)$$

and $K(u) = \frac{(1+u)^2}{4u}$.

Proof . From inequality (2.6), it follows that

$$\|\Phi^p(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \Phi^{-p}(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k)\| \leq (K(u^k))^p.$$

Lemma 2.3 leads to

$$\begin{pmatrix} K^p(u^k)I & \gamma_1 \\ \gamma_2 & K^p(u^k)I \end{pmatrix} \geq 0$$

and

$$\begin{pmatrix} K^p(u^k)I & \gamma_2 \\ \gamma_1 & K^p(u^k)I \end{pmatrix} \geq 0,$$

where

$$\gamma_1 = \Phi^p(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k)\Phi^{-p}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k)$$

and

$$\gamma_2 = \Phi^{-p}(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k)\Phi^p(A_1\sigma_2B_1, \dots, A_k\sigma_2B_k).$$

By summing up the two above inequalities, we obtain the following inequality:

$$\begin{pmatrix} 2(K^p(u^k))I & \gamma_1 + \gamma_2 \\ \gamma_2 + \gamma_1 & 2(K^p(u^k))I \end{pmatrix} \geq 0.$$

Again using Lemma 2.3, we obtain result. \square

Now, utilizing the continuous functional calculus, we show the following result.

Theorem 2.18. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ be such that $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$), σ_1 and σ_2 be two arbitrary matrix means between the arithmetic and harmonic means and $f, g : [m, M] \rightarrow [0, \infty)$ be two continuous functions such that g is non-zero, increasing and concave. Then, for every positive unital multilinear mapping $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_1(\mathbb{C})$

$$f(\Phi(A_1\sigma_1B_1, \dots, A_n\sigma_1B_n)) \leq \gamma g(\Phi(A_1\sigma_2B_1, \dots, A_n\sigma_2B_n))$$

where $\alpha_g = \frac{g(M)-g(m)}{M-m}$, $\beta_g = \frac{Mg(m)-mg(M)}{M-m}$, $u = \frac{M}{m}$, $K(u) = \frac{(1+u)^2}{4u}$ and

$$\gamma = \max \left\{ \frac{f(t)}{\alpha_g K^{-1}(u^k)t + \beta_g} : t \in [m, M] \right\}.$$

Proof . It follows from concavity of g that for $t \in [m, M]$

$$g(t) \geq \alpha_g t + \beta_g.$$

On the other hand, by our assumptions $m \leq \Phi(A_1\sigma_1B_1, \dots, A_n\sigma_1B_n) \leq M$. Applying monotonicity principle for operator functions and using lower Heinz inequality for(2.3), we get

$$\begin{aligned} g(\Phi(A_1\sigma_2B_1, \dots, A_n\sigma_2B_n)) &\geq \alpha_g \Phi(A_1\sigma_2B_1, \dots, A_n\sigma_2B_n) + \beta_g \\ &\geq \alpha_g K^{-1}(u^k)\Phi(A_1\sigma_1B_1, \dots, A_n\sigma_1B_n) + \beta_g \end{aligned} \tag{2.14}$$

where $K(u^k) = \frac{(1+u^k)^2}{4u^k}$. From $\alpha_g K^{-1}(u^k)t + \beta_g \neq 0$ for $t \in [m, M]$, it concludes that the function $\frac{f(t)}{\alpha_g K^{-1}(u^k)t + \beta_g}$ is continuous on the interval $[m, M]$. Now, we can put

$$\gamma = \max \left\{ \frac{f(t)}{\alpha_g K^{-1}(u^k)t + \beta_g} : t \in [m, M] \right\}.$$

Therefore, $\gamma(\alpha_g K^{-1}(u^k)t + \beta_g) \geq f(t)$. Again with the aid of monotonicity principle for operator functions, we get

$$\gamma(\alpha_g K^{-1}(u^k)\Phi(A_1\sigma_1B_1, \dots, A_n\sigma_1B_n) + \beta_g) \geq f(\Phi(A_1\sigma_1B_1, \dots, A_n\sigma_1B_n)). \tag{2.15}$$

Combining inequalities (2.14) and (2.15), we derive the desired result. \square

3 Some inequalities for dual and adjoint of matrix means

The authors in [11] introduced the adjoint and the dual of a matrix mean as follows: Let σ be an operator mean with representing function f . A matrix mean with representing function $f(t^{-1})^{-1}$ and $tf(t)^{-1}$ are called the adjoint and the dual of σ , respectively, and are denoted by σ^* and σ^\perp , respectively. Thus by the definition, we have

$$A\sigma^*B = (A^{-1}\sigma B^{-1})^{-1} \quad \text{and} \quad A\sigma^\perp B = (B^{-1}\sigma A^{-1})^{-1}.$$

The next Theorem is the key result of this section.

Theorem 3.1. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ such that $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$) with $0 < m < M$ and σ be an arbitrary matrix mean. Then

$$\Phi^2(A_1\sigma B_1, \dots, A_k\sigma B_k) \leq K^2(u^k)\Phi^2(A_1\sigma^* B_1 \cdots A_k\sigma^* B_k),$$

where σ^* is adjoint σ , $K(u) = \frac{(1+u)^2}{4u}$ and $u = \frac{M}{m}$.

Proof . Applying [4, Proposition 3.6],

$$\begin{aligned} & \Phi(A_1\sigma B_1, \dots, A_k\sigma B_k) + M^k m^k \Phi(A_1^{-1}\sigma B_1^{-1}, \dots, A_k^{-1}\sigma B_k^{-1}) \\ & \leq \Phi(A_1, \dots, A_k)\sigma\Phi(B_1, \dots, B_k) + M^k m^k (\Phi(A_1^{-1}, \dots, A_k^{-1})\sigma\Phi(B_1^{-1}, \dots, B_k^{-1})). \end{aligned}$$

By Lemma 2.5 and the subadditivity and the monotonicity property of the matrix mean σ , respectively, we derive

$$\Phi(A_1\sigma B_1, \dots, A_k\sigma B_k) + M^k m^k \Phi(A_1^{-1}\sigma B_1^{-1}, \dots, A_k^{-1}\sigma B_k^{-1}) \leq M^k + m^k. \quad (3.1)$$

Using Lemma 2.2(i), Lemma 2.1 and inequality (3.1) and applying a method similar to (2.3), the rest of proof is trivial. \square

As a consequence, we have the following result:

Remark 3.2. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ such that $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$), let σ be an arbitrary matrix mean and $p > 0$. Then,

$$\Phi^p(A_1\sigma B_1, \dots, A_k\sigma B_k) \leq \left(\frac{(1+u^k)^2}{4u^k} \right)^p \Phi^p(A_1\sigma^* B_1 \cdots A_k\sigma^* B_k)$$

for every positive unital multilinear mapping $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_n(\mathbb{C})$ and $u = \frac{M}{m}$. For $0 \leq p \leq 2$, by Theorem 3.1 and the Lowner-Heinz inequality, the result is clear. If $p > 2$, then using a similar method in Theorem 3.2 and using of Lemma 2.2(ii), we get the desired result.

Using the same proof as in Theorem 2.11, we have the following interesting Corollary.

Corollary 3.3. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ be such that $0 < m \leq A_i, B_i \leq M$ ($1 \leq i \leq k$) with $0 < m < M$, let σ_1 and σ_2 be two arbitrary matrix means between arbitrary matrix mean σ and its adjoint σ^* . Then, for every multilinear mapping $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_l(\mathbb{C})$,

$$\Phi^p(A_1\sigma_1 B_1, \dots, A_k\sigma_1 B_k) \leq K^p(u^k)\Phi^p(A_1\sigma_2 B_1, \dots, A_k\sigma_2 B_k), \quad p > 0 \quad (3.2)$$

where $u = \frac{M}{m}$ and $K(u) = \frac{(1+u)^2}{4u}$.

Remark 3.4. Taking $\sigma = \nabla$, then $\sigma^* = !$ and the inequality (3.2) becomes the inequality (2.5). Thus, this result generalize the obtained results in the first section.

Now, we present inequality (1.2) for the dual of a matrix mean.

Theorem 3.5. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ such that $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$) with $0 < m < M$ and σ be an arbitrary matrix mean. Then, for every multilinear mapping $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_l(\mathbb{C})$

$$\Phi^2(A_1\sigma B_1, \dots, A_k\sigma B_k) \leq K^2(u^k)\Phi^2(B_1\sigma^\perp A_1 \cdots B_k\sigma^\perp A_k),$$

where σ^\perp is the dual of σ , $u = \frac{M}{m}$ and $K(u) = \frac{(1+u)^2}{4u}$.

Proof . Notice that, by Lemma 2.2(i) and the inequality (3.1), we get

$$\|\Phi(A\sigma B, \dots, A\sigma B)M^k m^k \Phi^{-1}(B_1\sigma^\perp A_1, \dots, B_k\sigma^\perp A_k)\| \leq (M^k + m^k).$$

This shows the assertion as desired. \square

Using the same idea of the adjoint of a matrix mean, we can state:

Corollary 3.6. Let $A_i, B_i \in \mathcal{M}_n(\mathbb{C})$ be such that $0 < m \leq A_i, B_i \leq M$ ($i = 1, \dots, k$) with $0 < m < M$, let σ_1 and σ_2 be two arbitrary matrix means between arbitrary matrix mean σ and its dual σ^\perp . Then, for every multilinear mapping $\Phi : \mathcal{M}_n^k(\mathbb{C}) \rightarrow \mathcal{M}_l(\mathbb{C})$,

$$\Phi^p(A_1\sigma_1B_1, \dots, A_k\sigma_1B_k) \leq K^p(u^k)\Phi^p(B_1\sigma_2A_1, \dots, B_k\sigma_2A_k), \quad p > 0,$$

where $u = \frac{M}{m}$ and $K(u) = \frac{(1+u)^2}{4u}$.

Acknowledgements

The author would like to thank the Lorestan University.

References

- [1] T. Ando and X. Zhan, *Norm inequalities related to operator monotone functions*, Math. Ann. **315** (1999), 771–780.
- [2] M. Bakherad, *Refinements of a reversed AM-GM operator inequality*, Linear Multilinear Algeb. **64** (2016), no. 9, 1687–1695.
- [3] R. Bhatia and F. Kittaneh, *Notes on matrix arithmetic-geometric mean inequalities*, Linear Algebra Appl. **308** (2000), no. 1-3, 203–211.
- [4] M. Dehghani, M. Kian, and Y. Seo, *Developed matrix inequalities via positive multilinear mappings*, Linear Algebra Appl. **484** (2015), 63–85.
- [5] X. Fu and C. He, *Some operator inequalities for positive linear maps*, Linear Multilinear Algeb. **63** (2015), no. 3, 571–577.
- [6] X. Fu and D.T. Hoa, *On some inequalities with matrix means*, Linear Multilinear Algeb. **63**(2015), no. 12, 2373–2378.
- [7] T. Furuta, J. Mičić Hot, J. Pečarić, and Y. Seo, *Mond Pečarić Method in Operator Inequalities*, Element, Zagreb, 2005.
- [8] D.T. Hoa, D.T. H. Binh, and H.M. Toan, *On some inequalities with matrix means*, RIMS Kokyukoku, Kyotou Univ. **1893** (2014), no. 5, 67–71.
- [9] R.A. Horn and C.R. Johnson, *Topics in Matrix Analysis*, Cambridge University Press, 1991.
- [10] M. Kian and M. Dehghani, *Extension of the Kantorovich inequality for positive multilinear mappings*, Filomat **31** (2017), no. 20, 6473–6481.
- [11] F. Kubo and T. Ando, *Means of positive linear operators*, Math. Ann. **246** (1980), 205–224.
- [12] M. Lin, *Squaring a reverse AM-GM inequality*, Studia Math. **215** (2013), no. 2, 187–194.
- [13] J. Micic, J. Pecaric, and Y. Seo, *Complementary inequalities to inequalities of Jensen and Ando based on the Mond-Pecaric method*, Linear Algebra Appl. **318** (2000), 87–107.
- [14] L. Nasiri and M. Bakherad, *Improvements of some operator inequalities involving positive linear maps via the Kantorovich constant*, Houston J. Math. **45** (2019), no. 3, 815–830.
- [15] L. Nasiri and W. Liao, *The new reverses of Young type inequalities for numbers, operators and matrices*, Oper. Matrices **12** (2018), no. 4, 1063-1071.